
Breast Pathology


Elena Provenzano

46 yo female.


Hx Familial Adenomatous
Polyposis.


Lump R breast.





• • Case 1: B catenin





Fibromatosis

• • Fibromatosis

- Long intersecting fascicles of bland spindled cells
- Infiltrative edge invades around pre existing structures
- Nuclear β-catenin expression in 80% (but may also be seen in PT, spindle cell ca)
- Can be ER positive
- Negative for CD34, cytokeratins

• • Fibromatosis


- o DDx
- Scar look for foamy macrophages, haemosiderin, fat necrosis
- Spindle cell ca cytological atypia, epithelioid foci, DCIS, CK positivity
- Myofibroblastoma CD 34 positive


• • Fibromatosis


- Locally infiltrative benign neoplasm
- Local recurrence in 20-30%
- Reports of spontaneous regression
- Association with previous trauma including surgery, implants
- Can be seen in patients with FAP but majority in breast sporadic
- Activating mutations in β-catenin gene in 45%
- APC gene mutation or 5q loss in 33%


75 yo female.

Central L breast lump.









CK MNF 116


CK AE1/ AE3


Metaplastic Carcinoma Fibromatosis-like


• • Metaplastic carcinoma


- Low grade adenosquamous
- o Fibromatosis-like
- Spindle cell carcinomas
- Matrix producing carcinomas
- Squamous cell carcinomas

- 'All spindle cell lesions of the breast are metaplastic carcinoma until proven otherwise'
 - Sarah Pinder









Metaplastic carcinoma

- o Cam5.2 21% -52%
- AE1/AE3 79% -85%
- o EMA − 33% positive
- Broad spectrum CK 96% positive, 74% in >50% cells
- CK8/18 82-100%
- CK5 39%; CK14 33%
- CK7 17%
- \circ CD34 0%
- NEED TO USE A PANEL OF CYTOKERATINS INCLUDING BROAD SPECTRUM AND BASAL CYTOKERATINS
- STAINING CAN BE VERY FOCAL

Metaplastic carcinoma

	CK	p63	Nuclear B- catenin	CD34
Phyllodes tumour	- (rare focal)	- (rare focal)	+	+/-
Metaplastic carcinoma	+ (often focal)	+/-	-/+	-
Fibromatosis	-	-	+	-

- Focal CK positivity seen in PT MNF116 12%, CK7 28%, CK14 and Cam5.2 2%, AE1/3 8%. Associated with malignant subtype.
- Focal CK and p63 positivity identified in malignant PT (21% and 57% respectively);
- CD34 57% malignant PT positive, no staining in sarcomatoid carcinoma

Metaplastic carcinoma

Clues – associated NST carcinoma or DCIS


nuclear atypia or mitoses associated chronic inflammation


ER/ PgR/ HER2 negative


• • Metaplastic carcinoma

- Behaviour intermediate between carcinoma and sarcoma
- Intermediate risk of axillary LN metastases may relate to proportion of carcinomatous component
- Tendency to visceral metastases particularly lung
- Poor response to neoadjuvant chemotherapy
- Increased risk of local recurrence even with mastectomy – complete excision with good margins and RT important
- Poor survival: Median DFS 7 months (2-109/12)
- Median OS 18 months (4-109)


63 yo female. L breast lump.









ER







SMMHC

p63

Adenoid Cystic Carcinoma

Adenoid cystic carcinoma

Low grade lesion with cribriform architecture DDx:

- oLow grade DCIS
- Cribriform carcinoma
- Adenoid cystic carcinoma

•ER negative = Adenoid cystic carcinoma

Adenoid cystic carcinoma

- Salivary gland tumour also found in breast
- <1% of breast cancers</p>
- •Two cell populations;
- obasal/ myoepithelial cells CK5/6, CK14, p63
- oluminal/ epithelial cells CK7, CK8/18, CD117
- Triple negative with expression of basal markers
- oBUT older age group
- •Commonly grade 1 or 2
- oLow incidence nodal involvement up to 5%
- Excellent prognosis with 88-98% 5 year survival


Adenoid cystic carcinoma


- Distinct genetic profile to other basal breast cancers
- Less genetic instability, fewer copy number aberrations, intact BRCA1 pathway
- Characteristic translocation t(6:9) with formation of MYB
 NFIB fusion gene


35 yo female.

L breast lump.

M3U3 ? fibroadenoma









Carcinoma with Medullary Features

Carcinoma with medullary features

- Circumscribed tumour with pushing rather than infiltrating margin
- Interconnecting sheets of large, bizarre and pleomorphic carcinoma cells forming a syncytial network
- Prominent lymphocytic inflammatory cell infiltrate
- Usually ER/ HER2 negative
- Association with BRCA1 mutations





Basal Breast Cancers

- o 'Triple negative' ER, PgR and HER2 negative
- Express basal cytokeratins CK 5/6, CK14
- Express EGFR
- Distinct morphology high grade, central acellular zones, necrosis, high mitotic count
- Associated with BRCA1 mutations in young women
- Associated with worse prognosis and distant metastasis, particularly visceral metastases

Basal breast cancers

- Heterogeneous group of tumours
- o Grade 3 NST
- Medullary-like
- Metaplastic carcinomas

sarcomatoid


squamous


fibromatosis – like


low grade adenosquamous

- Adenoid cystic carcinoma
- SUBSET WITH A GOOD PROGNOSIS

81 yo female.
R breast mass
discovered on CT.









SMMHC

p63





Papilloma with florid usual type hyperplasia

Latest WHO Classification

- Intraductal papilloma
- o Intraductal papilloma with atypia/ DCIS
- Intraductal papillary carcinoma / Papillary DCIS
- Encapsulated papillary carcinoma
- Solid papillary carcinoma
- o Invasive papillary carcinoma

Papillary lesions


Classification dependent upon


- presence of cytological and architectural atypia
- presence of an internal and/ or external myoepithelial layer

Histopathological characteristics

	Intraductal Papilloma	Papilloma with ADH/ DCIS	Papillary DCIS	Encapsulated Papillary Carcinoma	Solid Papillary Carcinoma
Architecture	Broad, blunt fronds	Broad, blunt fronds	Slender fronds, sometimes branching	Slender fronds, sometimes branching	Solid with inconspicuous delicate fibro vascular septa
Epithelial Component	Heterogeneous +/- haphazard orientation: luminal cells, myoepithelial cells, usual hyperplasia, apocrine metaplasia	Focal areas with architecture and cytology of DCIS, usually low grade (monotonous cell pop'n, solid or cribriform spaces) Background heterogeneous	Entire lesion composed of cells with architecture and cytology of DCIS; usually low/ intermediate grade	Entire lesion composed of cells with architecture and cytology of DCIS; usually low/ intermediate	Entire lesion composed of cells with low grade cytology Spindle cell component Neuroendocrine diff'n common May show mucin production
Adjacent ducts	Usual hyperplasia	DCIS	DCIS	DCIS	DCIS

Atypical features





		MEC in Papillae	MEC at Periphery	
Papilloma		Present	Present	
Papilloma with atypia/DCIS		Present – may be absent/ reduced in atypical areas	Present	
Papillary DCIS		Absent	Present	
Encapsulated Papillary Ca		Absent	Absent	
Solid Papillary Ca		Absent	Present/ Absent	
Invasive Papillary Ca		Absent	Absent	

Collins LC, Histopathology 2008, 52; 20-29.

• • • IHC in diagnosis of papillary lesions

Two main settings


- Myoepithelial markers p63, SMMHC, calponin
- ER and basal CKs for epithelial proliferations to distinguish usual hyperplasia from ADH/ DCIS

• • • IHC – ER and basal CK's


- UDH is a polyclonal proliferation with a mosaic pattern of staining for ER and CK5/ CK14*
- ADH/ DCIS is a monoclonal proliferation with uniform strong staining for ER and negative staining for CK5/ CK14**

- * Apocrine proliferations are ER negative
- ** Basal DCIS is very rare in papillary lesions, but is ER negative and CK5/ CK14 positive with severe nuclear atypia
- ** This refers to epithelial cell staining, in papillomas with ADH/ DCIS myoepithelial cells will stain positively for basal CK's but appear as rare basally situated cells. With the mosaic pattern in UDH, 50-80% cells positive

• • IHC – Papilloma with UDH



• • IHC – Apocrine metaplasia



Papilloma with atypia/ DCIS


- Background benign papilloma with focal atypia/ DCIS
- DCIS usually low or intermediate grade with solid or cribriform architecture
- Myoepithelium may be reduced or absent in areas of atypia
- ?? Definition

Page >3mm = DCIS - used in WHO


Tavassoli <33% = atypia, 30-90% = DCIS

Schnitt – architectural features of DCIS = DCIS regardless of extent

IHC – Papilloma with atypia



• • • IHC – Papilloma with DCIS





• • IHC – myoepithelial markers


- p63 high sensitivity 99-100%
 low stromal cross reactivity (90% specificity)
- Calponin 93% sensitivity cross reactivity with myofibroblasts (79% specificity)
- SMA 88% sensitivity
 high cross reactivity with myofibroblasts
 (65% specificity)
- SMMHC high sensitivity, but some cross reactivity with myofibroblasts and vascular smooth muscle


Tse et al. J Clin Pathol 2009; 62:407-13.


• • IHC – myoepithelial markers












77 yo female.


L breast lump.


MMG normal.

Indeterminate lesion on US.



• • Case 6:





Solid papillary carcinoma – mixed in situ and invasive

Solid Papillary Carcinoma

- Previous edition subtype of EPC now separate entity
- Expansive cellular nodules with delicate fibrovascular cores that are inconspicuous at low power
- Neuroendocrine differentiation, spindle cell morphology and mucin production are common
- Myoepithelial cells may be present or absent surrounding the nodule


Solid Papillary Carcinoma


- Presence of outer myoepithelial cell layer, or rounded nodular masses without an infiltrative pattern are regarded as DCIS
- Rare variants with irregularly shaped islands arranged in a jigsaw pattern in a desmoplastic stroma, without a MEC layer on IHC = invasive solid papillary carcinoma


74 yo female.


L breast lump.


? malignant.


















Encapsulated papillary carcinoma

• • Encapsulated Papillary Ca

- Variant of papillary carcinoma characterised by fine fibrovascular cores covered by neoplastic cells of low to intermediate grade, surrounded by a thick fibrous capsule
- Lacks a myoepithelial layer
- Low incidence of lymph node metastasis (1-3%)
- Local recurrence rare (7%)
- Rare cases showing skeletal muscle invasion only if previous surgery
- Favourable prognosis with adequate local excision alone
- WHO stage as Tis unless frank invasive component
- Associated invasive carcinoma usually NST; present in 13-32% of cases
- For invasion requires infiltration beyond fibrous capsule
- Measure invasive component only

Epithelial displacement


- Common finding in resection specimens following CB
- Nests of epithelial cells lying within the biopsy tract – fibrosis, fat necrosis, haemosiderin laden macrophages
- In case of papillary carcinoma cells will be atypical
- Myoepithelial stains unhelpful
- Should not be called invasion




46 yo female.


Private screening MMG.


Distortion RUOQ.






• • Case 8: ER






• • Case 8: p63 and SMMHC



Invasive tubular and cribriform carcinoma with low grade DCIS

Tubular Carcinoma


- Composed of small, oval or rounded tubules lined by a single layer of epithelium – no myoepithelial layer
- Tumour cells are small and uniform with minimal nuclear atypia; cytoplasmic apical snouts may be present
- Connective tissue stroma is usually desmoplastic
- ER positive and HER2 negative
- Excellent prognosis





56 yo female.


Known Ca R breast.

Ca++ LUOQ on MMG M3.












Complex Sclerosing Lesion with focal LCIS

• • Radial scar/ CSL





• • Sclerosing adenosis





Sclerosing adenosis - SMMHC






• • • Radial Scar/ CSL

- B3 lesion
- Risk of associated malignancy is related to the presence of cytological atypia
 - Atypia present 28-44% upgraded
 - No atypia 4-6%
- Malignancy is usually in the form of low grade DCIS or low grade invasive carcinoma, particularly tubular carcinomas

• • • Case 10:

83 yo female. L breast lump.



• • Case 10 CK7 ER CK20 GATA3

• • • Case 10:

Invasive micropapillary carcinoma


Invasive Micropapillary Ca

- Reverse polarity with 'inside-out' tubules
- o 1-2% of invasive breast cancers
- Majority ER positive
- 10-30% HER2 positive may show incomplete pattern strong staining
- LVI and LN involvement common, however stage matched prognosis similar to NST


• • • Case 11:

50 yo female.
Suspicious lump L
breast and palpable
axillary lymph node.


• • • Case 11




• • • Case 11: ER




• • BUT.....pleural aspirate





• • • WT1



Pleural aspirate

Breast biopsy

• • • Case 11:

Metastatic Serous Papillary Carcinoma

Metastasis to Breast

- 0.2-1.3% of breast masses
- Presenting symptom in 30%
- Clinical history

88% previous history of malignancy 30% 1st site of met with no other disease 10% previous met, isolated breast disease 54% disseminated disease

Examination findings

77% mass, 85% unilateral, 53% solitary 15% skin/ subcutaneous nodules 2% inflammatory 1% diffuse parenchymal involvement

- Median survival 10 months
- Improved survival with surgery 15 vs 8 months
 Williams et al. Cancer 2007;110:731-7

Metastasis to Breast

- Histological findings:
 appearance not typical of breast primary, DCIS, elastosis
 and Ca++ all favour breast origin
- rarely clues such as pigment
- o Patterns:

discrete mass growth around ducts/ lobules (esp lymphomas) lymphangitis diffuse infiltration

Lee et al. J Clin Pathol 2007;60:1333-41

Metastasis to Breast

• Commonest sites:

melanoma
haematological malignancies
lung
ovary (serous papillary ca)

prostate, gastric, renal cell ca carcinoid tumours

Lee et al. J Clin Pathol 2007;60:1333-41

| Metastasis to Breast – IHC

Breast		CK7 +, CK20 – EMA 100% GCDFP 70% (23-45%)	
Serous Papillary Ca Ovary	WT1 95% Ca125 90% Mesothelin >90%	<10%, weak, focal 10-20% 3-15%, weak	
Lung	TTF1 – small cell, adenoca	small cell can be pos, primary breast rare	
Prostate	PSA PSAP CK7 & CK20 negative	15% positive 0% positive	
Gastric	ER negative CDX2 20-70% CK20 50% positive	Lobular ER positive	
Renal Cell Carcinoma	CD10 Renal cell ca marker 90%	5% positive 15% positive	
Carcinoid/ Neuroendocrine	ER negative CDX2/ CK20 – GI origin TTF1 – lung/thyroid	ER positive	


Breast Markers


Marker	Sensitivity	Specificity	Other Tumours	Comment
ER	85-90% primary, 50% mets		Gynae tract, thyroid, skin adenexal	Strong positive staining, weak may be non-sp
GCDFP- 15	10-70% (TNBC 10- 20%)	98-100%	Apocrine glands Rare ovarian and lung primaries	Apocrine differentiation Often focal
GATA3	>90% (TNBC 40- 70%)	70-93%	Urothelial ca Skin, salivary gland, kidney, SqCC	Nuclear Typically diffuse
Mamma globin	50-70% (TNBC 7- 23%)	90-100%	Salivary, skin adnexal, 5- 10% ovary/ endometrial	


• • • Case 12:


78 yo female.
R breast lump.
M5U5.

• • • Case 12









• • • Case 12:

Invasive lobular carcinoma – pleomorphic variant

• • • Pleomorphic Lobular Ca

- Growth pattern of lobular carcinoma but marked nuclear pleomorphism
- May have apocrine or histiocytoid morphology
- Majority GCDFP15 positive
- 75% ER positive (cf 96% classical ILC)
- 15-40% HER2 positive
- Worse survival compared with classical ILC


• • IHC - Ecadherin

- Ecadherin is a membrane bound protein involved in cell adhesion and forms part of the cadherin-catenin complex
- Loss of expression of Ecadherin is a feature of lobular carcinoma (LCIS and Invasive Lobular Carcinoma)
- BUT expression of Ecadherin is preserved in around 12% of ILC
- AND 25-50% of Invasive Ca NST (ductal) show reduced or absent expression of Ecadherin, especially high grade basal-like cancers
- Tumours are primarily classified as lobular or NST/ ductal on the basis of their morphology NOT Ecadherin staining
- Ecadherin IHC is most useful in CIS and for invasive cancers with a single file growth pattern where you are not sure if it is NST or lobular


• • IHC - Ecadherin

- Can get aberrant patterns of Ecadherin staining
- Partial fragmented membrane staining
- Cytoplasmic staining with perinuclear dot-like pattern
- Remember residual myoepithelial cells or ductal cells will be positive

Invasive lobular carcinoma




• • NST "with lobular features"



• • IHC – p120 catenin


- p120 catenin
- Cytoplasmic localisation in lobular lesions
- Membrane localisation in ductal/ NST lesions





Mastracci et al., Modern Pathology 2005, 18:741-51

• • • Case 13:


55 yo female.
WLE and Axillary
clearance 2010.
Now unresolving bruise
L breast.








• • • Case 13: CD34 and CD31





• • • Case 13:


Post Radiation Angiosarcoma

Post Radiation Angiosarcoma

- Originates in dermis, cf primary AS which arises within breast parenchyma
- Vascular proliferation with atypia; nuclear enlargement and hyperchromasia, necrosis and mitoses in high grade lesions
- Atypia may be minimal at periphery Ddx AVL
- AS poorly circumscribed, infiltrates subcutis
- CD31 and CD34 positive
- Epithelioid variant can be CK positive panel of markers
- Association with amplification of the MYC gene (55% of cases)
- Results in MYC protein overexpression IHC can be useful in differential diagnosis from AVL

Post RadiationAngiosarcoma

- Association with amplification of the MYC gene (55% of cases)
- Results in MYC protein overexpression IHC can be useful in differential diagnosis from AVL





Post Radiation Angiosarcoma


- o Two contexts:
 - Chest wall post radiotherapy
 - Upper limb following chronic lymphoedema post mastectomy
- o Rare: 1.27/1000 patients
- Latency 6-204 months; maximum risk at 5-10 yrs
- Locally aggressive tendency to multiple local recurrences before development of metastases
- Therapy surgical excision with clear margins
- Role for chemotherapy

• • • Case 14:

50 yo female.
Breast implants.
New lump R breast.











CK MNF CD 45

CD2









ALK CD 30

• • • Case 14:

Breast Implant Associated Anaplastic Large Cell Lymphoma (BIA-ALCL)

• • BIA-ALCL

- CD30 positive, ALK negative
- Variable expression of T cell markers; CD4 (80-84%), CD43 (80-88%), CD3 (30-46%), CD2 (30%)
- CD45 positive in 36%
- Negative for CD5, CD7, CD8 and CD15
- Up to 70% positive for EMA
- Majority of tested cases show monoclonal Tcell receptor gene rearrangements

• • BIA-ALCL


- Primary ALCL of the breast 3/100 million/ year
- 18.2 times risk with implants but still incredibly rare –
 1/500,00 women with implants
- BIA-ALCL arises within the implant capsule not the breast parenchymaCommonest presentation: late onset seroma causing breast swelling
- Excellent prognosis recommended management is removal of and and observe
- Patients who prcapsuleimplantesent with a mass / extension beyond the capsule may require more aggressive treatment with chemotherapy and / or radiotherapy


Breast Lymphoma


- Primary lymphoma of the breast 0.04-0.5% of breast tumours
- Conversely, 1-2% of extranodal lymphomas arise in the breast
- Commonest is diffuse large B cell lymphoma
- Next commonest are low grade B cell lymphomas;
 MALT and follicular lymphomas
- Rare Burkitt lymphoma, ALCL, peripheral T cell lymphomas


Case 17


Clinical history:
43 year old female
Diagnostic excision
following B4 core biopsy





















- Biphasic lesion with epithelial and myoepithelial components
- Mild cytological atypia
- Heterogeneous staining for ER
- Foci at periphery ? Stromal invasion
- o B4










• • WLE specimen



Final Diagnosis:

- Adenomyoepithelioma with atypia
- Areas at periphery ? Invasion
 => Emad for 2nd opinion
- P63 positivity throughout
- Lack of high grade atypia or other features of frank malignancy
- o Final diagnosis = atypical

• • Adenomyoepithelioma

- Biphasic tumour composed of a proliferation of myoepithelial cells around small epithelial spaces
- Resembles epithelial-myoepithelial carcinoma of the salivary gland
- Majority of patients post menopausal but can occur at any age; rare cases described in men
- Presents as a solitary mass lesion; may be history of rapid growth in a longstanding mass with malignant transformation
- Mammography circumscribed mass with lobulated outline similar in appearance to a fibroadenoma
- U/S solid or combined solid and cystic mass
- 1-8 cm in size; mean 2.5 cm

• • Adenopmyoepithelioma

- Circumscribed lesion with multilobulated architecture on low power, may have central sclerosis
- Lobulated, papillary, tubular or mixed patterns
- Papillary lesions with myoepithelial hyperplasia best regarded as papillomas
- Myoepithelial component predominates and may be spindled or epithelioid; latter often has clear cell appearance. Myoid and plasmacytoid variants also described.
- Epithelial component can show squamous, apocrine or sebaceous metaplasia
- Epithelial elements may be difficult to detect on light microscopy, particularly in spindle cell variant - IHC
- Lacks cytological atypia, mitoses rare (<3/10 HPF)

• • Adenomyoepithelioma

- Dual cell population demonstrated by IHC
- Epithelial component positive for CK7, CK8/18, EMA and ER
- Myoepithelial component positive for p63, SMMHC, calponin, actin, CK5/6, CK14. May stain with S100. Negative for desmin.

• • Adenomyoepithelioma

- Differential diagnosis includes papilloma, tubular adenoma, sclerosing adenosis, fibroadenoma
- Distinguishing feature is degree of presence of myoepithelial hyperplasia throughout the lesion
- Papillary lesions without nests, nodules or increased proportion of MEC classed as papillomas
- Ductal adenoma central location with surrounding fibrous capsule (no fibrous


MalignantAdenomyoepithelioma


- Atypical adenomyoepithelioma nuclear pleomorphism, increased mitoses (>4-5/10 HPF)
- Malignant adenomyoepithelioma requires presence of associated benign AME for diagnosis
- Malignant transformation can occur in epithelial component, myoepithelial component, or both
- Malignant features: nuclear pleomorphism, increased mitoses, necrosis, overgrowth of myoepithelial component, infiltrative growth pattern


- Two patterns:
- Typical adenomyoepithelioma that merges with an adjacent infiltrating carcinoma
- Epithelial invasive carcinoma NST, spindle cell or squamous metaplastic carcinoma, undifferentiated carcinoma
- Associated carcinoma often triple negative
- Myoepithelial carcinoma
- Epithelial-myoepithelial carcinoma similar to poorly differentiated salivary gland counterpart
- Lesion with the low power appearance of adenomyoepithelioma but on close examination has marked cytological atypia and mitotic activity

57 yo female. WLE for high grade DCIS


• • Case 20:









• • • Case 20: IHC





• • • Case 20: IHC





79 yo male. Left 'breast' excision.
















Myofibroblastoma


24 yo female. Excision biopsy B3 lesion R breast






SMMHC p63

ER



CK 5/6



Complex sclerosing lesion with usual hyperplasia


• • • Case 23

Encapsulated papillary carcinoma with associated invasive ca NST

• • • Case 23


Invasive size

Whole tumour size



Invasive size


W<u>hole tumou</u>r size



Multifocal


Invasive size


Whole tumour size




46 yo female.


Ulcerating lesion left nipple.








• • Case 24: IHC



• • • Case 24: IHC





• • • Case 24


Diagnosis:


Nipple adenoma with florid usual type hyperplasia

60 yo female. Right mastectomy.









Diagnosis:

Low grade adenosquamous ca with DCIS and LCIS